

U.S. APPLICATION NO. (IF KNOWN, SEE 37 CFR 1.53) 09/936543		INTERNATIONAL APPLICATION NO. PCT/EP00/01797		ATTORNEY'S DOCKET NUMBER 112843-031	
--	--	--	--	---	--

21. The following fees are submitted:				CALCULATIONS PTO USE ONLY	
BASIC NATIONAL FEE (37 CFR 1.492 (a) (1) - (5)) :					
<input type="checkbox"/>	Neither international preliminary examination fee (37 CFR 1.482) nor international search fee (37 CFR 1.445(a)(2)) paid to USPTO and International Search Report not prepared by the EPO or JPO			\$1,000.00	
<input checked="" type="checkbox"/>	International preliminary examination fee (37 CFR 1.482) not paid to USPTO but International Search Report prepared by the EPO or JPO			\$860.00	
<input type="checkbox"/>	International preliminary examination fee (37 CFR 1.482) not paid to USPTO but international search fee (37 CFR 1.445(a)(2)) paid to USPTO			\$710.00	
<input type="checkbox"/>	International preliminary examination fee paid to USPTO (37 CFR 1.482) but all claims did not satisfy provisions of PCT Article 33(1)-(4)			\$690.00	
<input type="checkbox"/>	International preliminary examination fee paid to USPTO (37 CFR 1.482) and all claims satisfied provisions of PCT Article 33(1)-(4)			\$100.00	
ENTER APPROPRIATE BASIC FEE AMOUNT =				\$860.00	
Surcharge of \$130.00 for furnishing the oath or declaration later than months from the earliest claimed priority date (37 CFR 1.492 (e)). <input type="checkbox"/> 20 <input type="checkbox"/> 30				\$0.00	
CLAIMS	NUMBER FILED	NUMBER EXTRA	RATE		
Total claims	23 - 20 =	3	x \$18.00	\$54.00	
Independent claims	6 - 3 =	3	x \$78.00	\$234.00	
Multiple Dependent Claims (check if applicable). <input type="checkbox"/>				\$0.00	
TOTAL OF ABOVE CALCULATIONS =				\$1,148.00	
Reduction of 1/2 for filing by small entity, if applicable. Verified Small Entity Statement must also be filed (Note 37 CFR 1.9, 1.27, 1.28) (check if applicable). <input type="checkbox"/>				\$0.00	
SUBTOTAL =				\$1,148.00	
Processing fee of \$130.00 for furnishing the English translation later than months from the earliest claimed priority date (37 CFR 1.492 (f)). <input type="checkbox"/> 20 <input type="checkbox"/> 30				\$0.00	
TOTAL NATIONAL FEE =				\$1,148.00	
Fee for recording the enclosed assignment (37 CFR 1.21(h)). The assignment must be accompanied by an appropriate cover sheet (37 CFR 3.28, 3.31) (check if applicable). <input type="checkbox"/>				\$0.00	
TOTAL FEES ENCLOSED =				\$1,148.00	
				Amount to be: refunded	\$
				charged	\$

☒ A check in the amount of **\$1,148.00** to cover the above fees is enclosed.


☐ Please charge my Deposit Account No. _____ in the amount of _____ to cover the above fees.
A duplicate copy of this sheet is enclosed.

☒ The Commissioner is hereby authorized to charge any fees which may be required, or credit any overpayment to Deposit Account No. **02-1818** A duplicate copy of this sheet is enclosed.

NOTE: Where an appropriate time limit under 37 CFR 1.494 or 1.495 has not been met, a petition to revive (37 CFR 1.137(a) or (b)) must be filed and granted to restore the application to pending status.

SEND ALL CORRESPONDENCE TO:

Robert M. Barrett, Esq.
 Bell, Boyd & Lloyd LLC
 P.O. Box 1135
 Chicago, Illinois 60690-1135
 Tel: (312) 807-4204


 SIGNATURE
 Robert M. Barrett
 NAME
 30,142
 REGISTRATION NUMBER
 September 10, 2001
 DATE

IN THE UNITED STATES PATENT AND TRADEMARK OFFICE

Applicant: Reniero et al.
Appl. No.: PCT/EP00/01797
Filed: March 2, 2000
Title: LACTIC ACID BACTERIA STRAINS CAPABLE OF PREVENTING
DIARRHOEA
Art Unit: Unknown
Examiner: Unknown
Docket No.: 112843-031

Assistant Commissioner for Patents
Washington, DC 20231

PRELIMINARY AMENDMENT

Sir:

Please amend the above-identified patent application as follows:

In the Claims:

Please amend Claims 3 and 6-11 as follows:

3. (Once Amended) The lactobacillus according to claim 1 which is selected from the group consisting of *Lactobacillus rhamnosus* and *Lactobacillus paracasei*.

6. (Once Amended) A method for preparing an ingestible support material comprising the step of using lactic acid bacterium belonging to the genus *Lactobacillus* capable of adhering to and essentially colonizing the intestinal mucosa and capable of preventing infection of intestinal epithelial cells by rotaviruses.

7. (Once Amended) The method according to claim 6, wherein the *Lactobacillus* strain is contained in the ingestible support material in an amount from about 10^5 cfu / g to about 10^{12} cfu / g support material.

8. (Once Amended) The method according to claim 6 wherein the ingestible support material is a food composition selected from the group consisting of milk, yogurt, curd, cheese, fermented milks, milk based fermented products, ice-creams, fermented cereal based products, milk based powders, and infant formulae.

9. (Once Amended) A method for the treatment of a disorder associated with diarrhoea comprising the step of administering to a patient suffering a disorder associated with diarrhoea lactic acid bacterium belonging to the genus *Lactobacillus* capable of adhering to and essentially colonizing the intestinal mucosa and capable of preventing infection of intestinal epithelial cells by rotaviruses.

10. (Once Amended) A pharmaceutical composition comprising a lactic acid bacterium belonging to the genus *Lactobacillus* capable of adhering to and essentially colonizing the intestinal mucosa and capable of preventing infection of intestinal epithelial cells by rotaviruses.

11. (Once Amended) The composition according to claim 10, which is selected from the group consisting of milk, yogurt, curd, cheese, fermented milks, milk based fermented products, ice-creams, fermented cereal based products, milk based powders, infant formulae, tablets, liquid bacterial suspensions, dried oral supplement, liquid oral supplement, dry tube feeding and liquid tube feeding.

Please add newly submitted claims 12-23 as follows:

12. The pharmaceutical composition according to claim 10 wherein the *Lactobacillus* is capable to grow in the presence of up to 0.4% bile salts.

13. The pharmaceutical composition according to claim 10 wherein the *Lactobacillus* is selected from the group consisting of *Lactobacillus rhamnosus* and *Lactobacillus paracasei*.

14. The pharmaceutical composition according to claim 13 wherein the *Lactobacillus* is *Lactobacillus paracasei*.

15. The pharmaceutical composition according to claim 13 wherein the *Lactobacillus paracasei* strain is *Lactobacillus paracasei* CNCM I-2116.

16. The method of claim 9 wherein the lactic acid bacterium comprises an ingestible support material.

17. The method according to claim 9 wherein the *Lactobacillus* strain is contained in the support material in an amount from about 10^5 cfu / g to about 10^{12} cfu / g support material.

18. A method for preventing a disorder associated with diarrhea comprising the steps of administering a lactic acid bacterium belonging to the genus *Lactobacillus* capable of adhering to and essentially colonizing the intestinal mucosa and capable of preventing infection of intestinal epithelial cells by rotaviruses to a patient at risk of diarrhea.

19. The method according to claim 18 wherein the lactic acid bacterium is part of a composition selected from the group consisting of milk, yogurt, curd, cheese, fermented milks, milk based fermented products, ice-creams, fermented cereal based products, milk based powders, and infant formula.

20. A food comprising a lactic acid bacterium belonging to the genus *Lactobacillus* capable of adhering to and essentially colonizing the intestinal mucosa and capable of preventing infection of intestinal epithelial cells by rotaviruses

21. The food of claim 20 which is selected from the group consisting of milk, yogurt, curd, cheese, fermented milks, milk based fermented products, ice-creams, fermented cereal based products, milk based powders, infant formulae, tablets, liquid bacteria suspensions, dried oral supplement, liquid oral supplement, dry tube feeding and liquid tube feeding.

22. The food of claim 20 wherein the Lactobacillus is capable of growing in the presence of up to 0.4% bile salts.

23. The food of claim 20 wherein the Lactobacillus is selected from the group consisting of Lactobacillus rhamnosus and Lactobacillus paracasei.

REMARKS

This Preliminary Amendment is submitted in the above-identified patent application. Pursuant to the Preliminary Amendment Claims 3 and 6-11 have been amended and newly submitted Claims 12-23 have been added. This Preliminary Amendment does not add new matter. Further, Applicants note that the Preliminary Amendment is not being made for purposes of patentability but, merely to comport the claims to U.S. practice and/or to add additional claims. Applicants do not intend to disclaim any subject matter in view of the amendments.

Attached hereto is a marked-up version of the changes made to the claims by the current amendment. The attached page is captioned "**Versions with Markings to Show Changes Made.**"

Respectfully submitted,

BELL, BOYD & LLOYD LLC

BY 

Robert M. Barrett
Reg. No. 30,142
P.O. Box 1135
Chicago, Illinois 60690-1135
Phone: (312) 807-4204

VERSION WITH MARKINGS TO SHOW CHANGES MADE

In the Claims:

Claims 3 and 6-11 have been amended as follows:

3. (Once Amended) The lactobacillus according to claim 1 [or claim 2], which is selected from the group consisting of *Lactobacillus rhamnosus* [or] and *Lactobacillus paracasei*.

6. (Once Amended) A method for preparing [use of a lactic acid bacterium according to any of the preceding claims for the preparation of] an ingestable support material comprising the step of using lactic acid bacterium belonging to the genus *Lactobacillus* capable of adhering to and essentially colonizing the intestinal mucosa and capable of preventing infection of intestinal epithelial cells by rotaviruses.

7. (Once Amended) The method [use] according to claim 6, wherein the *Lactobacillus* strain is contained in the ingestable support material in an amount from about 10^5 cfu / g to about 10^{12} cfu / g support material.

8. (Once Amended) The method [use] according to [any of the claims] claim 6 [or 7] wherein the ingestable support material is a food composition selected from the group consisting of milk, yogurt, curd, cheese, fermented milks, milk based fermented products, ice-creams, fermented cereal based products, milk based powders, and infant formulae.

9. (Once Amended) A method [The use according to any of the claims 5 to 7, wherein the support material used] for the treatment [and/or prophylaxis] of a disorder[s] associated with diarrhoea comprising the step of administering to a patient suffering a disorder associated with diarrhoea lactic acid bacterium belonging to the genus *Lactobacillus* capable of adhering to and essentially colonizing the intestinal mucosa and capable of preventing infection of intestinal epithelial cells by rotaviruses.

10. (Once Amended) A [Food or] pharmaceutical composition comprising a [containing] lactic acid bacterium belonging to the genus *Lactobacillus* capable of adhering to and essentially colonizing the intestinal mucosa and capable of preventing infection of intestinal epithelial cells by rotaviruses [at least once lactic acid bacterium according to any of the claims 1 to 4].

11. (Once Amended) The [composition] method according to claim [10,] 9 wherein the lactic acid bacteria is in a composition which is selected from the group consisting of [from] milk, yogurt, curd, cheese, fermented milks, milk based fermented products, ice-creams, fermented cereal based products, milk based powders, infant formulae, tablets, liquid bacterial suspensions, dried oral supplement, [wet] liquid oral supplement, dry tube feeding and [or] liquid [wet] tubefeeding.

Claims 12-23 were added.

LACTIC ACID BACTERIA STRAINS CAPABLE OF PREVENTING DIARRHOEA

The present invention pertains to novel microorganisms of the family Lactobacillaceae, especially to microorganisms of the genus Lactobacillus, that are useful in preventing or treating diarrhoea. In particular, the present invention relates to the use of said microorganisms for the preparation of an ingestible support and to a composition containing the same.

Organisms that produce lactic acid as a major metabolic component have been known for a long time. These bacteria may be found in milk or in milk processing factories, respectively, living or decaying plants but also in the intestine of man and animals. These microorganisms, summarized under the term "lactic acid bacteria", represent a rather inhomogeneous group and comprise e.g. the genera Lactococcus, Lactobacillus, Streptococcus, Bifidobacterium, Pediococcus etc..

Lactic acid bacteria have been utilized as fermenting agents for the preservations of food taking benefit of a low pH and the action of fermentation products generated during the fermentative activity thereof to inhibit the growth of spoilage bacteria. To this end, lactic acid bacteria have been used for preparing a variety of different foodstuff such as cheese, yogurt and other fermented dairy products from milk.

Quite recently lactic acid bacteria have attracted a great deal of attention in that some strains have been found to exhibit valuable properties to man and animals upon ingestion. In particular, specific strains of Lactobacillus or Bifidobacterium have been found to be able to colonize the intestinal mucosa and to assist in the maintenance of the well-being of man and animal.

In this respect, EP 0 768 375 discloses specific strains of the genus *Bifidobacterium*, that are capable to become implanted in the intestinal flora and may adhere to intestinal cells. These *Bifidobacteria* are reported to assist in immunomodulation, being capable to competitively exclude adhesion of pathogenic bacteria to intestinal cells, thus assisting in the maintenance of the individual's health.

During the last few years research has also focused on the potential use of lactic acid bacteria as probiotic agents. Probiotics are considered to be viable microbial preparations which promote the individual's health by preserving the natural microflora in the intestine. A microbial preparation may be commonly accepted as a probiotic in case the effectual microbes thereof and their mode of action are known. Probiotics are deemed to attach to the intestine's mucosa, colonize the intestinal tract and likewise prevent attachment of harmful microorganisms thereon. A crucial prerequisite for their action resides in that they have to reach the gut's mucosa in a proper and viable form and do not get destroyed in the upper part of the gastrointestinal tract, especially by the influence of the low pH prevailing in the stomach.

In this respect, WO 97/00078 discloses a specific strain, termed *Lactobacillus* GG (ATCC 53103), as such a probiotic. The microorganism is particularly employed in a method of preventing or treating food induced hypersensitivity reactions in that it is administered to a recipient together with a food material that has been subjected to a hydrolysis treatment with pepsin and/or trypsin. The *Lactobacillus* strain selected is described as exhibiting adhesive and colonizing properties and showing a protease enzyme system, so that the protein material contained in the foodstuff to be administered is further hydrolysed by means of proteases secreted by the specific *Lactobacillus* strain. The method discussed in this document shall eventually result in the uptake of protein material by the gut that does not show a substantial amount of allergenic material anymore.

Further, in EP 0 577 903 reference is made to the use of such lactic acid bacteria having the ability of replacing *Helicobacter pylori*, the acknowledged cause for the development of ulcer, in the preparation of a support intended for the therapeutic or prophylactic treatment of an ulcer associated with the action of *Helicobacter pylori*.

In knowledge of the valuable properties particular strains of lactic acid bacteria may provide there is a desire in the art for additional lactic acid bacterial strains that are beneficial to the well being of man and/or animal.

Consequently, a problem of the present invention is to provide additional bacterial strains that exhibit new properties beneficial for man and/or animals.

The above problem has been solved by providing novel microorganisms, namely lactic acid bacteria, belonging to the genus *Lactobacillus* having the traits of being capable to adhere to and essentially colonize the intestinal mucosa and to prevent infection of intestinal epithelial cells by rotaviruses.

According to a preferred embodiment the *Lactobacillus* strains are capable to grow in the presence of up to 0.4 % bile salts, so that they may easily pass the gastrointestinal tract and stay essentially active.

According to another preferred embodiment the lactic acid bacterium is selected from the group consisting of *Lactobacillus rhamnosus* or *Lactobacillus paracasei*, preferably *Lactobacillus paracasei*, and is more preferably *Lactobacillus paracasei* CNCM I-2116.

The microorganisms of the present invention have been shown to exhibit the following properties, they are gram positive, catalase negative, NH_3 form arginine negative and CO_2 production negative. They produce L(+) lactic acid, are capable to

grow in the presence of bile salts in a concentration of up to about 0.4 % and may essentially prevent infection of epithelial cells by rotaviruses.

The novel microorganisms may be used for the preparation of a variety of ingestible support materials, such as e.g. milk, yogurt, curd, fermented milks, milk based fermented products, fermented cereal based products, milk based powders, infant formulae and may be included in the support in an amount of from about 10^5 cfu / g to about 10^{11} cfu / g. For the purpose of the present invention the abbreviation cfu shall designate a "colony forming unit" that is defined as number of bacterial cells as revealed by microbiological counts on agar plates.

The present invention also provides a food or a pharmaceutical composition containing at least one of the Lactobacillus strains having the above traits.

For preparing a food composition according to the present invention at least one of the Lactobacillus strains according to the present invention is incorporated in a suitable support, in an amount of from about 10^5 cfu / g to about 10^{11} cfu / g, preferably from about 10^6 cfu / g to about 10^{10} cfu / g, more preferably from about 10^7 cfu / g to about 10^9 cfu / g.

In case of a pharmaceutical preparation the product may be prepared in forms of tablets, liquid bacterial suspensions, dried oral supplements, wet oral supplements, dry tube feeding or a wet tube feeding etc., with the amount of Lactobacillus strains to be incorporated therein being in the range of up to 10^{12} cfu / g, preferably from about 10^7 cfu / g to about 10^{11} cfu / g, more preferably from about 10^7 cfu / g to about 10^{10} cfu / g.

The activity of the novel microorganisms in the individual's intestine is naturally dose dependent. That is, the more the novel microorganisms are incorporated by means of ingesting the above food material or the pharmaceutical composition the higher the

protective and/or curing activity of the microorganisms. Since the novel microorganisms are not detrimental to mankind and animals and have eventually been isolated from baby feces a high amount thereof may be incorporated so that essentially a high proportion of the individual's intestine will be colonized by the novel microorganisms.

In the figures,

Fig. 1 shows a scheme illustrating the cell culture screening for assessing rotaviral protective properties of bacterial strains.

Fig. 2 shows the acidification of the *L. casei* strain CNCM I-2116 (termed ST11) in different growth media.

Fig. 3 shows the survival rate of the *L. casei* strain ST11 at 10 °C measured during 30 days.

Fig. 4 shows the mRNA pattern of IL-12 and IL-10 in mouse adherent cells derived from bone marrow after incubation of the cells with serial dilutions of ST11.

Fig. 5 shows the outcome of Th2 differentiation as resulting in decreased IL-4 production.

During the extensive studies leading to the present invention the inventors have investigated baby feces and isolated a variety of different bacterial strains therefrom. These strains were subsequently examined for their capability to prevent infection of epithelial cells with rotaviruses that are known to cause diarrhoea.

Several bacterial genera comprising *Lactobacillus*, *Lactococcus*, *Streptococcus* were screened for their rotavirus inhibitory properties. The tests for the inhibitory property

were essentially performed with three rotavirus serotypes representing the major etiological agents of human viral diarrhoea (serotypes G1, G3 and G4).

The various lactic acid bacteria were grown in a suitable medium, such as MRS, Hugo-Jago or M17 medium at temperatures of from about 30 to 40°C corresponding to their optimal growth temperature. After reaching stationary growth the bacteria were collected by centrifugation and resuspended in physiological NaCl solution. Between the different tests the bacterial cells were stored frozen (-20°C).

The various rotavirus stocks were prepared by infection of confluent cell monolayers. The rotaviruses were incubated before infection. The cells were infected with 20 tissue culture infectious doses.

For assessing anti-rotaviral properties two different protocols were applied. According to one protocol the various bacterial strains were examined for their direct interaction with the rotavirus while in the second protocol the bacteria were screened for those strains that interact with cellular rotavirus receptors.

The first protocol involved contacting the respective bacterial suspension each with a different rotavirus strain and incubating in suitable media. Subsequently, the virus-bacterium mixture was applied to a monolayer of cells of the human undifferentiated colon adenoma cells HT-29 and incubation was continued. Virus replication was then assayed.

The second protocol involved incubating the respective bacterial suspension first together with a monolayer of cells of the human undifferentiated colon adenoma cells HT-29 and adding the virus subsequently. After continued incubation virus replication was assayed.

Rotavirus replication may easily be assessed by histo-immunological staining of rotavirus proteins in infected cells.

A rotavirus inhibitory effect was attributed to a given bacterium when the number of infected cells was reduced by 90% in the cell culture inoculated with rotavirus plus the indicated bacteria in comparison with cells inoculated only with rotavirus.

Out of a total of 260 different bacterial strains primarily isolated merely 9 could be shown to essentially inhibit rotaviral replication. The different bacteria were ascertained to belong to the genus *Lactobacillus* subspecies *ramnosus* or *paracasei*. One strain, termed *Lactobacillus casei* ST11, that has been deposited in accordance with the Budapest Treaty and has received the deposit numbers NCC 2461 (I-2116), has been shown to be extremely effective in preventing infection of human cells by rotavirus. Moreover, this particular strain shows excellent growing properties as may be shown by acidification in different media. The strain also shows good performance as regards the survival rate during storage at low temperatures of about 10 °C, which makes it an excellent candidate for being included in food stuff or pharmaceutical compositions to be stored at refrigerator conditions.

In addition to the above finding it could also be shown that the strains surprisingly also exhibit anti-allergenic properties in that said strains have an impact on the synthesis of different immunological mediators.

It is generally acknowledged that humoral immune responses and allergic reactions are mediated by CD4⁺ T cells bearing the type 2 phenotype (Th2). Th2-cells are characterized by the production of high levels of interleukin 4 (IL-4), a cytokine required for the secretion of IgE, which is the major antibody class involved in allergic reactions.

The differentiation of Th2 cells is impaired by IFN- γ , a particular cytokine that arises from the mutually exclusive Th1 subset of CD4⁺ T cells. Said Th1 cells are in turn strongly induced by interleukin 12 (IL-12). In contrast thereto IL-10, another cytokine, has been shown to have a strong suppressing impact on the proliferation of Th1 cells and is therefore deemed to play a role in immunosuppressive mechanisms.

In summary, both IL-12 and IL-10 have strong modulatory effects on CD4⁺ T cell development by influencing the development of the Th1 subset. IL-12 is a key regulatory cytokine for the induction of Th1 differentiation and thus inhibits the generation of Th2 responses. A major pathway for inhibition of Th2 cells is therefore seen in the stimulation of IL-12 synthesis by accessory cells.

It is well known that some components of gram negative bacteria, such as LPS, induce high levels of IL-12 in adherent cells, such as macrophages and dendritic cells. Consistently, it has been found that gram negative bacteria can strongly bias CD4⁺ T cell differentiation towards the Th1 phenotype.

The microorganism ST11 as an example of the Lactobacillus strains of the present invention has been tested for a potential role in the induction of cytokines involved in the regulation of CD4⁺ T cell differentiation. In particular, the effect of ST11 on the phenotype of CD4⁺ T cells undergoing Th2 differentiation has been studied.

In this respect the capacity of ST11 to induce the synthesis of mRNA encoding these two regulatory cytokines in mouse adherent cells derived from bone marrow was compared with 4 other strains of Lactobacilli and with a control of gram negative bacteria (E.coli K12). The mRNA was measured by semi-quantitative RT-PCR after 6 hours of incubation of the cells with serial dilutions of bacteria ranging from 10⁹ to 10⁷ cfu/ml.

Although all strains of *Lactobacillus* could induce transcription of IL-12 mRNA to a certain degree, ST11 could be shown to be the strongest inducer, since as a strong PCR signal could be detected even at the lowest bacterial dose. In fact, the capacity of ST11 to induce IL-12 mRNA transcription was as strong as that of *E. coli*. Induction of IL-10 mRNA was in general weaker than for IL-12 mRNA, as only at higher bacterial doses a signal could be detected. Nevertheless, ST11 was the strongest inducer of IL-10 mRNA, as compared to the other lactobactilli and the *E. coli* control.

Thus, ST11 is deemed to be efficient in inducing immunoregulatory cytokines involved in CD4⁺ T cell differentiation. Its strong capacity to induce IL-12 makes it a candidate to inhibit Th2 responses and its measurable IL-10 induction may prevent inflammatory responses.

In addition to the above finding it was also determined whether ST11 had an inhibitory effect on CD4⁺ T cells undergoing Th2 differentiation and a positive effect on Th1 functions. A well established cell differentiation culture system was utilized, where precursor CD4⁺ T cells were polyclonally activated and modulated to undergo either Th1 or Th2 differentiation, depending on the type of co-stimuli provided in the culture medium. Th1/Th2 differentiation was induced during a 7-days primary culture, after which the cells were then restimulated for 2 days in a secondary culture containing medium alone and acquisition of a specific phenotype (Th1 or Th2) was assessed by measuring the types of cytokines produced in the supernatant (IFN- γ vs. IL-4).

It is known that precursor CD4⁺ T cells from mice of the BALB/c background preferentially differentiate to predominant Th2 phenotype (high IL-4, low IFN- γ in the 2ry culture supernatants) after activation under neutral conditions (medium alone in the 1ry culture). This phenotype could be completely reverted to a Th1 pattern

(high IFN- γ , low IL-4) upon addition of a blocking monoclonal antibody to IL-4 in the 1ry culture.

To investigate a potential role for ST11 on Th2 inhibition, purified precursor CD4⁺ T cells from BALB/c mice were activated in the presence of bone marrow adherent cells as accessory cells during the 1ry culture. These cells were co-cultured either in medium alone, or in the presence of 1 mg/ml LPS, or 10⁸ cfu/ml ST11, or 10⁸ cfu/ml of another Lactobacillus. After this time, the cells were washed and CD4⁺ T cells were purified once again and restimulated in the 2ry culture in medium alone. Cytokines produced by the differentiated CD4⁺ T cells were measured after 2 days. As expected, cells differentiated in the presence of medium alone displayed a dominant Th2 phenotype. Addition of ST11 to the 1ry cultures strongly modulated the outcome of Th2 differentiation, as it resulted in an 8-fold decrease in IL-4 production. This inhibition was of similar magnitude as that observed in cultures derived from cells differentiated in the presence of LPS. In contrast, the other Lactobacillus strain had no measurable impact on IL-4 levels. Interestingly, IFN- γ levels were not increased upon addition of ST11 in the 1ry cultures.

In summary, ST11 specifically impaired IL-4 production by CD4⁺ T cells undergoing Th2 differentiation, but did not significantly increase IFN- γ secretion. The fact that ST11 does not increase IFN- γ production may be due to its capacity to induce IL-10 with the consequence that it may keep a low inflammatory impact despite its anti-Th2 activity.

In consequence, it could be shown that ST11 is a Lactobacillus strain that has a good anti-Th2 profile which makes it an excellent candidate for its use as a bacterium with anti-allergic, probiotic activity.

The present invention will now be described by way of example.

Media and solutions:

MRS (Difco),

Hugo-Jago (Tryptone Difco 30 g/l, Yeast Extract Difco 10 g/l, Lactose Difco 5 g/l, KH_2PO_4 5 g/l, Beef Extract Difco 2 g/l, agar Difco 2 g/l)

M17 (Difco)

M199 (Seromed)

Ringer solution (Oxoid)

PBS (NaCl 8g/l, KCl 0.2 g/l, Na_2HPO_4 1.15 g/l, KH_2PO_4 0.2 g/l)

Tryptose phosphate broth (Flow)

Trypsin-EDTA solution (Seromed)

Human rotavirus Wa (G1 serotype) and simian rotavirus SA-11 (G3 serotype) were obtained from P.A. Offit, Children's Hospital of Philadelphia, U.S.A. The DS-1xRRV reassortant virus was obtained from A. Kapikian, NIH Bethesda, U.S.A. The serotype 4 human rotavirus Hocht was obtained from P. Bachmann, University of Munich, Germany.

Example 1**Isolation of lactic acid bacteria from baby feces**

Fresh feces were harvested from diapers of 16 healthy babies 15 to 27 days old. 1 g of fresh feces was placed under anaerobic conditions for transportation to the laboratory and microbiological analyses were run within 2 hours from sampling by serial dilutions in Ringer solution and plating on selective media. MRS agar plus antibiotics (phosphomycin 80 $\mu\text{g/ml}$, sulfamethoxazole 93 $\mu\text{g/ml}$, trimethoprim 5 $\mu\text{g/ml}$) incubated at 37°C for 48 hours was used to isolate lactic acid bacteria. Colonies were randomly picked up and purified. Physiological and genetic characterisation was performed on the isolates.

Example 2

Testing of strains in cell culture for anti-rotaviral activity

Several bacterial genera comprising *Lactobacillus*, *Lactococcus*, *Streptococcus* were selected and tested for members which showed anti-rotavirus activity in the cell culture inhibition test. The genus *Lactococcus* was represented by a single species (*Lc. lactis*) consisting of two subspecies (*Lc. lactis* supsp. *lactis* and *cremoris*). A total of 30 strains were tested. The *Streptococcus* genus was represented by a single species (*S. thermophilus*) with 45 strains. The *Leuconostoc* and *Propionibacterium* genus were only represented by a single species (6 strains), while the *Enterococcus* and *Staphylococcus* genus was represented by two species each and a total of 17 strains.

In total, 260 bacterial strains were tested for rotavirus inhibitory activity.

1st protocol:

30 µl of bacterial suspension (containing at average 3×10^6 bacteria) were mixed with 70 µl M199 medium supplemented with 10% tryptose phosphate broth (Flow) and 5% trypsin-EDTA solution (Seromed) (1 : 4 diluted in the case of HT-29 cells) and 100 µl of virus in supplemented M199 medium. The virus-bacterium mixture was incubated for 1 hour at 4°C and for 1 hour at 37°C. Cells of the human undifferentiated colon adenoma cells HT-29 growing as a confluent monolayer in 96-well microtiter plates were washed three times with phosphate-buffered saline (PBS ; pH 7.2). The virus-bacterium mixture was applied to the cells and the microtiter plates were incubated for 18 h in a CO₂ incubator (Heraeus). Virus replication was assayed as described below.

2nd protocol:

30 µl of the bacterial suspension (supra) were mixed with 70 µl M199 medium supplemented with 10% tryptose phosphate broth (Flow) and 5% trypsin-EDTA

solution (Seromed) (1 : 4 diluted in the case of HT-29 cells) and applied directly on the cells in the microtiter plates. After one hour incubation at 37°C 100 µl of virus in supplemented M199 medium was added to the cells in the microtiter plates. The incubation was continued for 18 h in a CO₂ incubator (Heraeus). Virus replication was assayed as described below.

The rotavirus replication was assessed by histo-immunological staining of rotavirus proteins in infected cells as described hereafter.

One day after infection, the cell culture medium was discarded from the microtiter plates and the cells were fixed with absolute ethanol for 10 min. Ethanol was discarded, and the plates were washed three times in PBS buffer. Then 50 µl of an anti-rotavirus serum (mainly directed against VP6 protein), produced in rabbits (obtained from the ISREC University of Lausanne) and diluted 1 :2000 in PBS was added to each well and incubated for 1 h at 37°C with a cover slip to prevent desiccation of the wells. The antiserum was discarded afterwards and the plates were washed three times with PBS. Then 50 µl of anti-rabbit immunoglobulin G (IgG) antiserum produced in goats and coupled to peroxidase (GAR-IgG-PO; Nordic) were added at a dilution of 1 : 500 in PBS to each well and the plates were incubated for 1 hour at 37 °C. The serum was discarded and the plates were once again washed three times with PBS. Then 100 µl of the following substrate mixture was added to each well : 10 ml of 0.05 M Tris-hydrochloride (pH 7.8), 1 ml of H₂O₂ (30% suprapur, diluted 1 :600 in H₂O ; Merck) and 200 µl of 3-amino-9-ethylcarbazole (0.1 g/10 ml of ethanol stored in 200 µl aliquots at -80 °C ; A-5754 ; Sigma). The plates were incubated for at least 30 min at room temperature. The substrate was discarded and the wells were filled with 200 µl of H₂O to stop the reaction. Infected cell foci were counted with an inverted microscope (Diavert ; Leitz).

Only very few bacterial strains interacted with rotaviruses. Merely 9 out of the 260 bacterial cells primarily selected inhibited rotavirus replication in at least one

protocol. *Lactobacillus paracasei* NCC 2461 (ST11) showed an extremely high activity against Serotype 1 Rotavirus, Serotype 3 rotavirus SA-11 and Serotype 4 rotavirus Hocht.

Example 3

Properties of ST11

ST 11 has been subjected to incubation in simulated gastric juice. The simulated gastric juice was prepared by suspending pepsin (3 g/l) in sterile saline (0.5% w/v) and adjusting the pH to 2.0 and pH 3, respectively, with concentrated HCl. ST 11 has been grown in varying amounts in the above media and the resistance of the microorganisms has been determined.

The results are summarized in table I below .

Table I

pH	cfu/ml at T 0	cfu at T 1min	Cfu at T 15	cfu at T 30	cfu at T 60
2.0	2.0×10^9	1.8×10^9	1.2×10^9	3.7×10^8	7.0×10^3
3.0	2.0×10^9	1.9×10^9	1.7×10^9	1.7×10^9	8.4×10^8

ST11 has the following properties as defined according to methods disclosed in the genera of lactic acid bacteria, Ed. B.J.B. Wood and W.H. Holzappel, Blackie A&P.

- gram positive,
- catalase negative,
- NH_3 form arginine negative
- CO_2 production negative
- production of L(+) lactic acid
- growth in the presence of bile salts in a concentration of up to about 0.4 %.

Example 4**Growth of ST 11 under different conditions**

ST11 was incubated at 37 °C in tomato based medium (4% tomato powder rehydrated in distilled water) supplemented with sucrose (0, 0.5, 1 or 2%) or soya peptone (0.5%) or glucose (0.5%) for different periods of time. The results are shown in fig. 2.

ST11 was further added in an amount of 2.5 % to a medium composed of rice flour (3%), wheat flour (2%) and sucrose (3%) and incubated at 37°C until a pH of 4.4 was reached. After cooling the product was packed with or without addition of vit.C and stored at 10°C.

Example 5**Induction of IL-12 and IL10- mRNA synthesis in mouse adherent cells by ST11**

Bone marrow cells were isolated from the femur and tibia of 8 week-old specific pathogen-free C57BL/6 mice and were incubated at a concentration of 2×10^6 cells/ml in RPMI medium (Gibco) containing 10% fetal bovine serum, 1 mM L-Glutamine, 12 mM Hepes, 0.05 mM 2-mercaptoethanol, 100 U/ml penicillin and 100 µg/ml streptomycin (all reagents from Gibco) for 12 hours at 37°C in a 5% CO₂ atmosphere. Non-adherent cells were discarded by 3 consecutive washes with warm culture medium and the remaining adherent cells were harvested and incubated at a concentration of 10^6 cells/ml for 6 hours in the presence or absence of bacteria. It was previously determined that 6 hours represents an optimal time-point for cytokine mRNA synthesis by mouse adherent cells in response to LPS. Bacteria were added at different concentrations ranging from 10^9 to 10^7 cfu/ml. Bacteria were grown and stored as indicated above (page 6).

At the end of the 6-hour culture period, cells were isolated by centrifugation and lysed using the TRIzol reagent kit (GibcoBRL, Cat. No. 15596-018) following the manufacturer's instructions. Total RNA was isolated by isopropanol precipitation and reverse-transcribed into cDNA for 90 min at 42°C using 200 U reverse transcriptase (Superscript II, BRL) in a 40- μ l reaction volume containing 200 mM Tris pH 8.3, 25 mM KCl, 1 μ g/ml oligo d(T)₁₅ (Boehringer Mannheim), 1 mM DTT (Boehringer Mannheim), 4 mM of each dNTP (Boehringer Mannheim) and 40 U/ml Rnasin (Promega). PCR primers and conditions were used as already described in Kopf et al. (Journal of Experimental Medicine 1996 Sep 1;184(3):1127-36). Amounts of cDNA were normalized within the samples using primers specific for a house-keeping gene (β -2-microglobulin). PCR products were separated on a 2% agarose gel and bands were analyzed under UV.

As shown in Figure 4, ST11 showed the strongest induction of IL-12 and IL-10 mRNA, which was comparable to levels observed with the positive control (*E. coli*). Differences are best seen at the lowest bacteria concentrations (10⁷ cfu/ml).

Example 6

Suppression of IL-4 synthesis by ST11

CD4⁺ T cells were purified from the spleen of specific pathogen-free BALB/c mice using the MiniMACS kit from Miltenyi Biotec (Cat. No. 492-01). The CD4⁺ T cells were cultured at a concentration of 2x10⁵ cells/ml in RPMI medium containing 10% fetal bovine serum, 1 mM L-Glutamine, 12 mM Hepes, 0.05 mM 2-mercaptoethanol, 100 U/ml penicillin and 100 μ g/ml streptomycin and activated during one week by cross-linking with plate-bound monoclonal antibodies to CD3 (clone 2C11) and CD28 (clone 37.51, both antibodies from Pharmingen). During this 1ry culture, the CD4⁺ T cells were co-cultured with bone marrow adherent cells (isolated as described above) as accessory cells and with 10⁸ cfu/ml ST11, or 10⁸ cfu/ml La1, or 1 mg/ml LPS, or medium alone. After this time, the cells were washed and CD4⁺ T cells were

- 17 -

purified once again using MiniMACS kit technology and restimulated in a 2ry culture containing medium alone. Cytokines produced by the differentiated CD4⁺ T cells were measured in the supernatants after 2 days using sandwich ELISA (kits from Endogen and Pharmingen).

Results are shown in Fig. 5. Cells differentiated in the presence of medium alone displayed a dominant Th2 phenotype characterized by high levels of IL-4. Addition of ST11 to the 1ry cultures strongly modulated the outcome of Th2 differentiation, as it resulted in an 8-fold decrease in IL-4 production. This inhibition was of similar magnitude as that observed in cultures derived from cells differentiated in the presence of LPS. In contrast thereto, the other Lactobacillus strain had no measurable impact on IL-4 levels. Interestingly, IFN- γ levels were not increased upon addition of ST11 in the 1ry cultures.

As may be seen from the above the strains of the present invention may be well prepared for the production of a food and/or a pharmaceutical carrier taking advantage of the valuable properties of the microorganisms.

Example 7

The ST11 strain was tested in a clinical trial in a community on the outskirts of Guatemala City on its ability to influence the transmission and experience of acute rainy-season diarrhoeal disease experienced by most of the children in this area. A total of 203 children, aged 35 to 70 months were enrolled in the study and received a target dose of 10¹⁰ viable organisms (ST11) or none (placebo) over a feeding period of 29 days. The children selected for both sample and placebo, respectively, had the typical deficits in weight-for-age and height-for-age characteristic due to undernutrition.

Before initiating the feeding trial in preschool children, a safety evaluation was conducted based on *in vitro* and *in vivo* studies. In vitro studies showed antibiotic resistance pattern similar to those of other lactobacilli used in food applications, and no potential for forming

biogenic amines, for degrading mucin and for deconjugating bile salts. In a placebo-controlled clinical study involving 42 adult volunteers, ST11 was well tolerated and did not induce any adverse effects, among the potential manifestations monitored, such as flatulence, number of stools per day and stool consistency; the levels of acute-phase proteins in serum did not raise any concern with respect to a potential inflammatory reaction.

The samples and placebo's have been packed in sachets at the Nestlé Product Technology Center manufacturing facility in Konolfingen, Switzerland, and shipped refrigerated to Guatemala. Each 10 g sachet consisted of a chocolate flavored vehicle and either 0.2 g of ST11 (10^{10} cfu) or, in case of the placebo, 0.2 g milk powder. The chocolate flavored vehicle consisted of cocoa powder, sugar, soy lecithin, vanillin, and cinnamon. The sachets were stored at 4° to 6°C until two hours before use. Before use the sachet had to be dissolved in 100 ml water provided by Nestlé, which was free of any bacterial contamination.

According to the protocol applied diarrhoea was defined as the occurrence of three or more liquid or unformed fecal evacuation during a period of 24 hrs. A diarrhoeal episode was defined as an event that presented the evidence of diarrhoea (3 diarrhoeal evacuations over 24 hrs). Its total duration in hrs was calculated from the moment of the first of the three index stools to the appearance of the first formed stool or a period 24 h without any defecation. For a child to have a "new" episode, 48 hours had to have elapsed the termination of the prior episode. If not, it was considered a continuation of the same episode and the total duration was then used for evaluation. A case was a child who experienced one or more documented episodes of diarrhoea through the 29-day observation period. Intensity of a diarrhoeal episode was based on the total number of loose stools produced. The elements of severity of the episode embraced the presence of blood, mucus or pus in stools, along with symptoms of fever and vomiting. An intensity of 7 stools per 24 h or the need for intervention by a health professional at a clinic, health center or hospital also classified an episode as severe.

When a diarrhoeal episode was diagnosed through the surveillance system, a specimen of a diarrhoeal stool was collected for microscopic examination and culture to identify potential

- 19 -

etiological pathogens for that episode. The specimen was diagnosed for rotavirus antigen, Giardia, and *E. histolytica*, in case the sample was dysenteric, and for bacterial pathogens including Shigella, Salmonella, Aeromonas, Plesiomonas shigelloides, *E. Coli* and possibly *V. cholerae*.

During the period of investigation product samples were collected to examine the viability of the microorganisms included during the period of administration. It could be shown that the microorganism stayed viable in the sachets during the entire study so that also at the end of the study the sachets were capable to convey 10^{10} viable microorganism upon reconstitution with water.

The study revealed that the sample containing the probiotic microorganism could decrease the occurrence of diarrhoea in contrast to the control group (placebo) by about 30 %. Yet, also the control group already exhibited a decreased number of diarrhoeal occurrence over the normal population receiving none of the samples or placebo, respectively. This latter finding may in part be explained on the basis of the children receiving additional valuable nutrition and contamination free water. However, since the study was performed in the field it may clearly be derived that ST11 can surely reduce the occurrence of diarrhoea in vivo.

TRAITE DE BUDAPEST ~~OR LA RECONNAISSANCE~~
INTERNATIONALE DU DEPOT DES MICRO-ORGANISMES
AUX FINS DE LA PROCEDURE EN MATIERE DE BREVETS

FORMULE INTERNATIONALE

DESTINATAIRE :

RECEPISSE EN CAS DE DEPOT INITIAL,
délivré en vertu de la règle 7.1 par
l'AUTORITE DE DEPOT INTERNATIONALE
identifiée au bas de cette page

SOCIETE DES PRODUITS NESTLE S.A.
Patents department
Avenue Nestlé 55
CH-1800 Vevey

NOM ET ADRESSE
DU DEPOSANT

I. IDENTIFICATION DU MICRO-ORGANISME

Référence d'identification donnée par le DEPOSANT :

NCC 2461Numéro d'ordre attribué par
l'AUTORITE DE DEPOT INTERNATIONALE :**I - 2116**

II. DESCRIPTION SCIENTIFIQUE ET/OU DESIGNATION TAXONOMIQUE PROPOSEE

Le micro-organisme identifié sous chiffre I était accompagné :



d'une description scientifique



d'une désignation taxonomique proposée

(Cocher ce qui convient)

III. RECEPTION ET ACCEPTATION

La présente autorité de dépôt internationale accepte le micro-organisme identifié sous
chiffre I, qu'elle a reçu le **12 JANVIER 1999** (date du dépôt initial)¹

IV. RECEPTION D'UNE REQUETE EN CONVERSION

La présente autorité de dépôt internationale a reçu le micro-organisme identifié sous
chiffre I le (date du dépôt initial)
et a reçu une requête en conversion du dépôt initial en dépôt conforme au Traité de
Budapest le (date de réception de la requête en conversion)

V. AUTORITE DE DEPOT INTERNATIONALE

Nom :

CNCM
Collection Nationale
de Cultures de Microorganismes

Adresse :

INSTITUT PASTEUR
28, Rue du Docteur Roux
F-75724 PARIS CEDEX 15

Signature(s) de la (des) personne(s)
compétente(s) pour représenter l'autorité
de dépôt internationale ou de l'(des)
employé(s) autorisé(s) : **Mme Y. CERISIER**
Directeur Administratif de la CNCM

Date : **Paris, le 12 février 1999**

¹ En cas d'application de la règle 6.4.d), cette date est la date à laquelle le statut
d'autorité de dépôt internationale a été acquis.

Claims

1. Lactic acid bacterium belonging to the genus *Lactobacillus* capable of adhering to and essentially colonizing the intestinal mucosa, capable of preventing infection of intestinal epithelial cells by rotaviruses and capable to grow in the presence of up to 0.4 % bile salts.

2. The *Lactobacillus* according to claim 1, which is selected from the group consisting of *Lactobacillus rhamnosus* or *Lactobacillus paracasei*.

3. The *Lactobacillus* according to claim 2, which is *Lactobacillus paracasei*.

4. The *Lactobacillus paracasei* strain according to claim 3, which is *Lactobacillus paracasei* CNCM I-2116.

5. Use of a lactic acid bacterium according to any of the preceding claims for the preparation of a ingestable support material.

6. The use according to claim 5, wherein the *Lactobacillus* strain is contained in the support material in an amount from about 10^5 cfu / g to about 10^{12} cfu / g support material.

7. The use according to any of the claims 5 or 6, wherein the support material is a food composition selected from milk, yogurt, curd, cheese, fermented milks, milk based fermented products, ice-creams, fermented cereal based products, milk based powders, infant formulae.

8. The use according to any of the claims 5 to 6, wherein the support material is used for the treatment and/or prophylaxis of disorders associated with diarrhea.

9. Food or pharmaceutical composition containing at least one lactic acid bacterium according to any of the claims 1 to 3.

- 5 10. The composition according to claim 9, which is selected from milk, yogurt, curd, cheese, fermented milks, milk based fermented products, ice-creams, fermented cereal based products, milk based powders, infant formulae, tablets, liquid bacterial suspensions, dried oral supplement, wet oral supplement, dry tube feeding or wet tubefeeding.

10

11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65
66
67
68
69
70
71
72
73
74
75
76
77
78
79
80
81
82
83
84
85
86
87
88
89
90
91
92
93
94
95
96
97
98
99
100
101
102
103
104
105
106
107
108
109
110
111
112
113
114
115
116
117
118
119
120
121
122
123
124
125
126
127
128
129
130
131
132
133
134
135
136
137
138
139
140
141
142
143
144
145
146
147
148
149
150
151
152
153
154
155
156
157
158
159
160
161
162
163
164
165
166
167
168
169
170
171
172
173
174
175
176
177
178
179
180
181
182
183
184
185
186
187
188
189
190
191
192
193
194
195
196
197
198
199
200
201
202
203
204
205
206
207
208
209
210
211
212
213
214
215
216
217
218
219
220
221
222
223
224
225
226
227
228
229
230
231
232
233
234
235
236
237
238
239
240
241
242
243
244
245
246
247
248
249
250
251
252
253
254
255
256
257
258
259
260
261
262
263
264
265
266
267
268
269
270
271
272
273
274
275
276
277
278
279
280
281
282
283
284
285
286
287
288
289
290
291
292
293
294
295
296
297
298
299
300
301
302
303
304
305
306
307
308
309
310
311
312
313
314
315
316
317
318
319
320
321
322
323
324
325
326
327
328
329
330
331
332
333
334
335
336
337
338
339
340
341
342
343
344
345
346
347
348
349
350
351
352
353
354
355
356
357
358
359
360
361
362
363
364
365
366
367
368
369
370
371
372
373
374
375
376
377
378
379
380
381
382
383
384
385
386
387
388
389
390
391
392
393
394
395
396
397
398
399
400
401
402
403
404
405
406
407
408
409
410
411
412
413
414
415
416
417
418
419
420
421
422
423
424
425
426
427
428
429
430
431
432
433
434
435
436
437
438
439
440
441
442
443
444
445
446
447
448
449
450
451
452
453
454
455
456
457
458
459
460
461
462
463
464
465
466
467
468
469
470
471
472
473
474
475
476
477
478
479
480
481
482
483
484
485
486
487
488
489
490
491
492
493
494
495
496
497
498
499
500
501
502
503
504
505
506
507
508
509
510
511
512
513
514
515
516
517
518
519
520
521
522
523
524
525
526
527
528
529
530
531
532
533
534
535
536
537
538
539
540
541
542
543
544
545
546
547
548
549
550
551
552
553
554
555
556
557
558
559
560
561
562
563
564
565
566
567
568
569
570
571
572
573
574
575
576
577
578
579
580
581
582
583
584
585
586
587
588
589
590
591
592
593
594
595
596
597
598
599
600
601
602
603
604
605
606
607
608
609
610
611
612
613
614
615
616
617
618
619
620
621
622
623
624
625
626
627
628
629
630
631
632
633
634
635
636
637
638
639
640
641
642
643
644
645
646
647
648
649
650
651
652
653
654
655
656
657
658
659
660
661
662
663
664
665
666
667
668
669
670
671
672
673
674
675
676
677
678
679
680
681
682
683
684
685
686
687
688
689
690
691
692
693
694
695
696
697
698
699
700
701
702
703
704
705
706
707
708
709
710
711
712
713
714
715
716
717
718
719
720
721
722
723
724
725
726
727
728
729
730
731
732
733
734
735
736
737
738
739
740
741
742
743
744
745
746
747
748
749
750
751
752
753
754
755
756
757
758
759
760
761
762
763
764
765
766
767
768
769
770
771
772
773
774
775
776
777
778
779
780
781
782
783
784
785
786
787
788
789
790
791
792
793
794
795
796
797
798
799
800
801
802
803
804
805
806
807
808
809
810
811
812
813
814
815
816
817
818
819
820
821
822
823
824
825
826
827
828
829
830
831
832
833
834
835
836
837
838
839
840
841
842
843
844
845
846
847
848
849
850
851
852
853
854
855
856
857
858
859
860
861
862
863
864
865
866
867
868
869
870
871
872
873
874
875
876
877
878
879
880
881
882
883
884
885
886
887
888
889
890
891
892
893
894
895
896
897
898
899
900
901
902
903
904
905
906
907
908
909
910
911
912
913
914
915
916
917
918
919
920
921
922
923
924
925
926
927
928
929
930
931
932
933
934
935
936
937
938
939
940
941
942
943
944
945
946
947
948
949
950
951
952
953
954
955
956
957
958
959
960
961
962
963
964
965
966
967
968
969
970
971
972
973
974
975
976
977
978
979
980
981
982
983
984
985
986
987
988
989
990
991
992
993
994
995
996
997
998
999
1000
1001
1002
1003
1004
1005
1006
1007
1008
1009
1010
1011
1012
1013
1014
1015
1016
1017
1018
1019
1020
1021
1022
1023
1024
1025
1026
1027
1028
1029
1030
1031
1032
1033
1034
1035
1036
1037
1038
1039
1040
1041
1042
1043
1044
1045
1046
1047
1048
1049
1050
1051
1052
1053
1054
1055
1056
1057
1058
1059
1060
1061
1062
1063
1064
1065
1066
1067
1068
1069
1070
1071
1072
1073
1074
1075
1076
1077
1078
1079
1080
1081
1082
1083
1084
1085
1086
1087
1088
1089
1090
1091
1092
1093
1094
1095
1096
1097
1098
1099
1100
1101
1102
1103
1104
1105
1106
1107
1108
1109
1110
1111
1112
1113
1114
1115
1116
1117
1118
1119
1120
1121
1122
1123
1124
1125
1126
1127
1128
1129
1130
1131
1132
1133
1134
1135
1136
1137
1138
1139
1140
1141
1142
1143
1144
1145
1146
1147
1148
1149
1150
1151
1152
1153
1154
1155
1156
1157
1158
1159
1160
1161
1162
1163
1164
1165
1166
1167
1168
1169
1170
1171
1172
1173
1174
1175
1176
1177
1178
1179
1180
1181
1182
1183
1184
1185
1186
1187
1188
1189
1190
1191
1192
1193
1194
1195
1196
1197
1198
1199
1200
1201
1202
1203
1204
1205
1206
1207
1208
1209
1210
1211
1212
1213
1214
1215
1216
1217
1218
1219
1220
1221
1222
1223
1224
1225
1226
1227
1228
1229
1230
1231
1232
1233
1234
1235
1236
1237
1238
1239
1240
1241
1242
1243
1244
1245
1246
1247
1248
1249
1250
1251
1252
1253
1254
1255
1256
1257
1258
1259
1260
1261
1262
1263
1264
1265
1266
1267
1268
1269
1270
1271
1272
1273
1274
1275
1276
1277
1278
1279
1280
1281
1282
1283
1284
1285
1286
1287
1288
1289
1290
1291
1292
1293
1294
1295
1296
1297
1298
1299
1300
1301
1302
1303
1304
1305
1306
1307
1308
1309
1310
1311
1312
1313
1314
1315
1316
1317
1318
1319
1320
1321
1322
1323
1324
1325
1326
1327
1328
1329
1330
1331
1332
1333
1334
1335
1336
1337
1338
1339
1340
1341
1342
1343
1344
1345
1346
1347
1348
1349
1350
1351
1352
1353
1354
1355
1356
1357
1358
1359
1360
1361
1362
1363
1364
1365
1366
1367
1368
1369
1370
1371
1372
1373
1374
1375
1376
1377
1378
1379
1380
1381
1382
1383
1384
1385
1386
1387
1388
1389
1390
1391
1392
1393
1394
1395
1396
1397
1398
1399
1400
1401
1402
1403
1404
1405
1406
1407
1408
1409
1410
1411
1412
1413
1414
1415
1416
1417
1418
1419
1420
1421
1422
1423
1424
1425
1426
1427
1428
1429
1430
1431
1432
1433
1434
1435
1436
1437
1438
1439
1440
1441
1442
1443
1444
1445
1446
1447
1448
1449
1450
1451
1452
1453
1454
1455
1456
1457
1458
1459
1460
1461
1462
1463
1464
1465
1466
1467
1468
1469
1470
1471
1472
1473
1474
1475
1476
1477
1478
1479
1480
1481
1482
1483
1484
1485
1486
1487
1488
1489
1490
1491
1492
1493
1494
1495
1496
1497
1498
1499
1500
1501
1502
1503
1504
1505
1506
1507
1508
1509
1510
1511
1512
1513
1514
1515
1516
1517
1518
1519
1520
1521
1522
1523
1524
1525
1526
1527
1528
1529
1530
1531
1532
1533
1534
1535
1536
1537
1538
1539
1540
1541
1542
1543
1544
1545
1546
1547
1548
1549
1550
1551
1552
1553
1554
1555
1556
1557
1558
1559
1560
1561
1562
1563
1564
1565
1566
1567
1568
1569
1570
1571
1572
1573
1574
1575
1576
1577
1578
1579
1580
1581
1582
1583
1584
1585
1586
1587
1588
1589
1590
1591
1592
1593
1594
1595
1596
1597
1598
1599
1600
1601
1602
1603
1604
1605
1606
1607
1608
1609
1610
1611
1612
1613
1614
1615
1616
1617
1618
1619
1620
1621
1622
1623
1624
1625
1626
1627
1628
1629
1630
1631
1632
1633
1634
1635
1636
1637
1638
1639
1640
1641
1642
1643
1644
1645
1646
1647
1648
1649
1650
1651
1652
1653
1654
1655
1656
1657
1658
1659
1660
1661
1662
1663
1664
1665
1666
1667
1668
1669
1670
1671
1672
1673
1674
1675
1676
1677
1678
1679
1680
1681
1682
1683
1684
1685
1686
1687
1688
1689
1690
1691
1692
1693
1694
1695
1696
1697
1698
1699
1700
1701
1702
1703
1704
1705
1706
1707
1708
1709
1710
1711
1712
1713
1714
1715
1716
1717
1718
1719
1720
1721
1722
1723
1724
1725
1726
1727
1728
1729
1730
1731
1732
1733
1734
1735
1736
1737
1738
1739
1740
1741
1742
1743
1744
1745
1746
1747
1748
1749
1750
1751
1752
1753
1754
1755
1756
1757
1758
1759
1760
1761
1762
1763
1764
1765
1766
1767
1768
1769
1770
1771
1772
1773
1774
1775
1776
1777
1778
1779
1780
1781
1782
1783
1784
1785
1786
1787
1788
1789
1790
1791
1792
1793
1794
1795
1796
1797
1798
1799
1800
1801
1802
1803
1804
1805
1806
1807
1808
1809
1810
1811
1812
1813
1814
1815
1816
1817
1818
1819
1820
1821
1822
1823
1824
1825
1826
1827
1828
1829
1830
1831
1832
1833
1834
1835
1836
1837
1838
1839
1840
1841
1842
1843
1844
1845
1846
1847
1848
1849
1850
1851
1852
1853
1854
1855
1856
1857
1858
1859
1860
1861
1862
1863
1864
1865
1866
1867
1868
1869
1870
1871
1872
1873
1874
1875
1876
1877
1878
1879
1880
1881
1882
1883
1884
1885
1886
1887
1888
1889
1890
1891
1892
1893
1894
1895
1896
1897
1898
1899
1900
1901
1902
1903
1904
1905
1906
1907
1908
1909
1910
1911
1912
1913
1914
1915
1916
1917
1918
1919
1920
1921
1922
1923
1924
1925
1926
1927
1928
1929
1930
1931
1932
1933
1934
1935
1936
1937
1938
1939
1940
1941
1942
1943
1944
1945
1946
1947
1948
1949
1950
1951
1952
1953
1954
1955
1956
1957
1958
1959
1960
1961
1962
1963
1964
1965
1966
1967
1968
1969
1970
1971
1972
1973
1974
1975
1976
1977
1978
1979
1980
1981
1982
1983
1984
1985
1986
1987
1988
1989
1990
1991
1992
1993
1994
1995
1996
1997
1998
1999
2000
2001
2002
2003
2004
2005
2006
2007
2008
2009
2010
2011
2012
2013
2014
2015
2016
2017
2018
2019
2020
2021
2022
2023
2024
2025
2026
2027
2028
2029
2030
2031
2032
2033
2034
2035
2036
2037
2038
2039
2040
2041
2042
2043
2044
2045
2046
2047
2048
2049
2050
2051
2052
2053
2054
2055
2056
2057
2058
2059
2060
2061
2062
2063
2064
2065
2066
2067
2068
2069
2070
2071
2072
2073
2074
2075
2076
2077
2078
2079
2080
2081
2082
2083
2084
2085
2086
2087
2088
2089
2090
2091
2092
2093
2094
2095
2096
2097
2098
2099
2100
2101
2102
2103
2104
2105
2106
2107
2108
2109
2110
2111
2112
2113
2114
2115
2116
2117
2118
2119
2120
2121
2122
2123
2124
2125
2126
2127
2128
2129
2130
2131
2132
2133
2134
2135
2136
2137
2138
2139
2140
2141
2142
2143
2144
2145
2146
2147
2148
2149
2150
2151
2152
2153
2154
2155
2156
2157
2158
2159
2160
2161
2162
2163
2164
2165
2166
2167
2168
2169
2170
2171
2172
2173
2174
2175
2176
2177
2178
2179
2180
2181
2182
2183
2184
2185
2186
2187
2188
2189
2190
2191
2192
2193
2194
2195
2196
2197
2198
2199
2200
2201
2202
22





Fig. 1


2/5

Fig. 2

Acidification of *L. casei* ST11 in different growth media

Survival in cereal drink

Fig. 3




HDPE: High density poly ethylene

PS: polystyrene

Vit C: vitamin C

4/5

Fig. 4 Induction of cytokine mRNA by LAB in mouse macrophages



5/5

CYTOKINE RELEASE IN IVD 980916
(ELISA 981103 & 980925)




Fig. 5

Docket No.
112843-031

Declaration and Power of Attorney For Patent Application

English Language Declaration

As a below named inventor, I hereby declare that:

My residence, post office address and citizenship are as stated below next to my name,

I believe I am the original, first and sole inventor (if only one name is listed below) or an original, first and joint inventor (if plural names are listed below) of the subject matter which is claimed and for which a patent is sought on the invention entitled
LACTIC ACID BACTERIA STRAINS CAPABLE OF PREVENTING DIARRHOEA

the specification of which

(check one)

☐ is attached hereto.

☒ was filed on 2 March 2000 as United States Application No. or PCT International Application Number PCT/EP00/01797
and was amended on February 28, 2001 and September 10, 2001
(if applicable)

I hereby state that I have reviewed and understand the contents of the above identified specification, including the claims, as amended by any amendment referred to above.

I acknowledge the duty to disclose to the United States Patent and Trademark Office all information known to me to be material to patentability as defined in Title 37, Code of Federal Regulations, Section 1.56.

I hereby claim foreign priority benefits under Title 35, United States Code, Section 119(a)-(d) or Section 365(b) of any foreign application(s) for patent or inventor's certificate, or Section 365(a) of any PCT International application which designated at least one country other than the United States, listed below and have also identified below, by checking the box, any foreign application for patent or inventor's certificate or PCT International application having a filing date before that of the application on which priority is claimed.

Prior Foreign Application(s)	Priority	Not Claimed
<u>PCT/EP00/01797</u>	<u>2 March 2000</u>	<input type="checkbox"/>
(Number)	(Day/Month/Year Filed)	
<u>99104924.8</u>	<u>11 March 1999</u>	<input type="checkbox"/>
(Number)	(Day/Month/Year Filed)	
		<input type="checkbox"/>
(Number)	(Day/Month/Year Filed)	

I hereby claim the benefit under 35 U.S.C. Section 119(e) of any United States provisional application(s) listed below:

(Application Serial No.)

(Filing Date)

(Application Serial No.)

(Filing Date)

(Application Serial No.)

(Filing Date)

I hereby claim the benefit under 35 U. S. C. Section 120 of any United States application(s), or Section 365(c) of any PCT International application designating the United States, listed below and, insofar as the subject matter of each of the claims of this application is not disclosed in the prior United States or PCT International application in the manner provided by the first paragraph of 35 U.S.C. Section 112, I acknowledge the duty to disclose to the United States Patent and Trademark Office all information known to me to be material to patentability as defined in Title 37, C. F. R., Section 1.56 which became available between the filing date of the prior application and the national or PCT International filing date of this application:

(Application Serial No.)

(Filing Date)

(Status)
(patented, pending, abandoned)

(Application Serial No.)

(Filing Date)

(Status)
(patented, pending, abandoned)

(Application Serial No.)

(Filing Date)

(Status)
(patented, pending, abandoned)

I hereby declare that all statements made herein of my own knowledge are true and that all statements made on information and belief are believed to be true; and further that these statements were made with the knowledge that willful false statements and the like so made are punishable by fine or imprisonment, or both, under Section 1001 of Title 18 of the United States Code and that such willful false statements may jeopardize the validity of the application or any patent issued thereon.

POWER OF ATTORNEY: As a named inventor, I hereby appoint the following attorney(s) and/or agent(s) to prosecute this application and transact all business in the Patent and Trademark Office connected therewith. (list name and registration number)



29174

PATENT & TRADEMARK OFFICE

Send Correspondence to: **Robert M. Barrett**
Bell, Boyd & Lloyd LLC
P.O. Box 1135
Chicago, Illinois 60690-1135

Direct Telephone Calls to: (name and telephone number)
Robert M. Barrett (312) 807-4204

Full name of sole or first inventor Roberto Reniero	1-00
Sole or first inventor's signature <i>[Signature]</i>	Date 7/11/01
Residence Le Mont-Pelerin, Switzerland	CHX
Citizenship Italy	
Post Office Address 24, chemin Paudille	
CH-1801 Le Mont-Pelerin, Switzerland	

Full name of second inventor, if any Harald Bruessow	2-00
Second inventor's signature <i>[Signature]</i>	Date 25-10-01
Residence La Tour de Peilz, Switzerland	CHX
Citizenship Germany	
Post Office Address Chemin de la Chaumeny 13	
CH-1814 La Tour de Peilz, Switzerland	

Full name of third inventor, if any Florence Rochat	3-00
Third inventor's signature <i>Florence Rochat</i>	Date 26.10.01
Residence Montreux, Switzerland	CHX
Citizenship Switzerland	
Post Office Address Quartier des Tilleuls 6	
CH-1820 Montreux, Switzerland	

Full name of fourth inventor, if any Thierry Von Der Weid	4-00
Fourth inventor's signature <i>Thierry Von Der Weid</i>	Date 05.11.01
Residence Paudex, Switzerland	CHX
Citizenship Switzerland	
Post Office Address Rte de la Bordinette 10	
CH-1094 Paudex, Switzerland	

Full name of fifth inventor, if any Stephanie Blum-Sperisen	5-00
Fifth inventor's signature <i>Stephanie Blum-Sperisen</i>	Date 26-10-01
Residence Lausanne, Switzerland	CHX
Citizenship Germany	
Post Office Address Av. des Mousquines 13	
CH-1005 Lausanne, Switzerland	

Full name of sixth inventor, if any	
Sixth inventor's signature	Date
Residence	
Citizenship	
Post Office Address	